

# Kinks and Time Machines

Andrew Chamblin, G.W. Gibbons, Alan R. Steif

*Department of Applied Mathematics and Theoretical Physics,  
Silver Street, Cambridge CB3 9EW, England.*

We show that it is not possible to smooth out the metric on the Deutsch-Politzer time machine to obtain an everywhere non-singular asymptotically flat Lorentzian metric.

There has been interest recently in asymptotically flat spacetimes containing closed timelike curves (CTCs). One widely discussed example of such a spacetime (sometimes referred to as the Deutsch-Politzer spacetime) is constructed as follows [1, 2]. One takes ordinary  $n$ -dimensional Minkowski spacetime with inertial coordinates  $(t, \mathbf{x})$  and deletes two spatial balls, i.e., one removes the points

$$t = 0, |\mathbf{x}| < 1$$

and

$$t = 1, |\mathbf{x}| < 1.$$

One now identifies the upper side of the earlier ball with the lower side of the later ball and the upper side of the later ball with the lower side of the earlier point. That is, one identifies points  $t = \epsilon, |\mathbf{x}| \leq 1$  with points  $t = 1 - \epsilon, |\mathbf{x}| \leq 1$  and points  $t = -\epsilon, |\mathbf{x}| \leq 1$  with points  $t = 1 + \epsilon, |\mathbf{x}| \leq 1$  having the same  $\mathbf{x}$  values, where  $\epsilon$  is an infinitesimal positive quantity. The metric is, as we shall formally demonstrate later, singular at the branch points  $t = 0, t = 1, |\mathbf{x}| = 1$ . It is clear that as a manifold, we can smooth out the singularities at  $t = 0, |\mathbf{x}| = 1$ , to obtain a smooth manifold with the topology of  $\mathbb{R}^n$  with a handle attached, i.e., which is homeomorphic to  $S^1 \times S^{n-1} - \{\text{pt}\}$ , where the point corresponds to infinity. It is less clear whether one can smooth out the spacetime metric to obtain a smooth time-orientable Lorentzian spacetime with an everywhere non-singular Lorentz

metric,  $g$ , which is asymptotically flat. In fact one cannot: no matter how one tries, the metric must always contain singularities. The point of this note is to give a simple proof of this fact in the case that  $n$  is even. This covers the two most interesting cases;  $n = 4$ , the physical case, and  $n = 2$ , which provides a simple toy model.

Our proof makes use of the properties of a *gravitational kink* which we have developed previously for dealing with topological problems involving Lorentzian metrics [3, 4, 5]. We begin by surrounding the region of CTCs,  $0 < t < 1$ ,  $|\mathbf{x}| < 1$ , with a large  $(n - 1)$ -sphere,  $S_\infty$ ,

$$t^2 + |\mathbf{x}|^2 = R^2$$

where  $R$  is much larger than the size of the time machine. We now have a compact manifold,  $M$ , with boundary  $\partial M \cong S^{n-1}$ .

The necessary and sufficient condition that any compact manifold,  $M$  with boundary  $\partial M$  admit a Lorentz metric  $g_L$  is that

$$\chi[M] = \text{kink}(\partial M; g_L) \tag{1}$$

where  $\chi[M]$  is the Euler characteristic of  $M$  and  $\text{kink}(\partial M; g_L)$  is the kink number of the Lorentz metric  $g_L$  with respect to the boundary  $\partial M$  [3]. The definition of the kink number  $\text{kink}(\partial M; g_L)$  is as follows. One introduces on  $M$  an auxillary Riemannian metric  $g_R$  and then diagonalises  $g_L$  with respect to  $g_R$ . Because  $g_L$  is assumed to be time-orientable, the field of eigenvectors with negative eigenvalue provides an everywhere non-vanishing vector field  $\mathbf{V}$ , which may be assumed to be normalised to have unit length with respect to the Riemannian metric  $g_R$ . The kink number  $\text{kink}(\partial M; g_L)$  is now the algebraic number of times that  $\mathbf{V}$  coincides with the inward pointing unit normal  $\mathbf{n}$  of  $\partial M$ . More precisely,  $\mathbf{V}$  and  $\mathbf{n}$  provide two global sections of the bundle  $S(\partial M)$  of unit 4-vectors over  $\partial M$ . Then two sections intersect generically in an isolated number of points which may be assigned a sign depending upon the orientations of the sections. One then counts the number of points with regard to sign. The orientation conventions are such that if one reverses the sign of the normal, then the sign of the kink number changes. Reversing the sign of the vector field  $\mathbf{V}$ , however, leaves the kink number unchanged.

In two spacetime dimensions, one may characterise the kink number in terms of the more familiar winding number. Let  $\theta$  be the angle through

which  $\mathbf{V}$  must be rotated counter-clockwise to coincide with  $\mathbf{n}$ . The kink number is the total change in  $\frac{\theta}{2\pi}$  as the boundary  $\partial M$  is traversed with the inward normal  $\mathbf{n}$  pointing left. Changing the sign of  $\mathbf{V}$  will merely change  $\theta$  to  $\theta + \pi$  everywhere on the boundary and thus leave invariant the total change in  $\theta$  and hence the kink number. However, changing the sign of  $\mathbf{n}$  will reverse the kink number because now the boundary must be traversed in the opposite direction (see Figure 1).

Applying equation (1) to ordinary Minkowski spacetime inside the large sphere  $S_\infty$  (or calculating directly) shows that

$$\text{kink}(S_\infty; g_L) = +1. \quad (2)$$

However, one has

$$\chi(S^1 \times S^{n-1} - \{\text{pt}\}) = (-1)^{n-1}. \quad (3)$$

Thus equation (1) cannot be satisfied for even  $n$ . More generally, we could consider a spacetime with  $g$  time machines. This would have the topology of the connected sum, denoted by  $\#$ , of  $\mathbb{R}^n$  with  $g$  handles. One has

$$\chi(\mathbb{R}^n \# S^1 \times S^{n-1} \# S^1 \times S^{n-1} \dots \# S^1 \times S^{n-1}, S_\infty) = 1 - g(1 + (-1)^n) \quad (4)$$

which again contradicts equation (1). We note in passing (see [5]) that these results also go through in the non-time-orientable case.

We now show directly that the branch points  $|\mathbf{x}| = 1$  are singular in the two-dimensional case. Consider a curve enclosing one of the branch points, say  $x = 1$ . As in the previous discussion, we calculate the winding number of the timelike vector field  $\frac{\partial}{\partial t}$  with respect to the inward normal to the curve. Since the curve winds first around the point  $t = 1$ ,  $|\mathbf{x}| = 1$  and then around  $t = 0$ ,  $|\mathbf{x}| = 1$ , one sees that the kink number is two. However, the kink number which is obtained upon encircling a non-singular point in Minkowski space is unity. Therefore, the spacetime is not locally Minkowskian at the branch point. This argument is readily generalised to higher dimensions.

It is perhaps worth emphasizing that the manifold  $S^1 \times S^{n-1} - \{\text{pt}\}$  does in fact admit a non-singular Lorentz metric. For example, take the vector field which winds around the  $S^1$  factor. This Lorentz metric however has kink number  $= -1$  and so cannot be asymptotically flat. This difference may be traced back to the fact that the kink number changes sign on reversal of the normal (see Figure 3).

We have shown (at least for even  $n$ ) that one cannot construct an asymptotically flat Deutsch-Politzer type time machine with an everywhere time-orientable non-singular Lorentz metric. For  $n = 2$ , it does not seem that one can construct a non-trivial non-singular model at all since all one is allowed is to add on more handles, but by equation (4), this makes things worse. In the case of four-dimensional spacetimes, it is clear that one can construct asymptotically flat time machines, but these must be considerably more complicated than the Deutsch-Politzer model.

From the physical point of view, it is not clear to us how seriously one should worry about the singularities in the Deutsch-Politzer spacetime. For the purposes of discussing conceptual problems involving CTCs, the view has been expressed to us that one may always impose boundary conditions at these singularities that prevent information from them affecting such issues as the unitarity of quantum field theory on these backgrounds. What is clear to us is that the gravitational kink concept is useful in detecting the presence of such singularities. In this respect, the situation resembles attempts to model topology change using a trouser-leg spacetime – i.e., one diffeomorphic to  $S^n$  with balls removed and with no spacelike boundary components. Equation (1) shows that no non-singular Lorentz metric is possible on the trouser-leg spacetime. If  $n = 4$ , one can find a spacetime with three spacelike boundary components – take  $\mathbb{CP}^2$  and remove three four-balls for example. However, the resulting spacetime does not admit a spinor structure. The results of [6] show that if  $n = 4$ , no such Lorentz cobordism admits a spinor structure. On the other hand, there will exist such cobordisms which will admit a globally defined Cliffordian *pin* structure (see [5]). Put another way, by allowing our cobordisms to be non-orientable (see [5]), we can have arbitrary *spacelike* topology change while preserving the global existence of a non-vanishing Lorentz metric and a ‘pinor’ bundle (which we can use to quantise the Dirac equation).

Of course, one could take the view that one must go beyond globally defined Lorentzian metrics and consider singular metrics if one wishes to adhere rigidly to the idea that a single spacetime metric is involved in topology change. Perhaps the least drastic modification of the globally defined Lorentzian metric concept is that of a metric with regions of different spacetime signature.

## ACKNOWLEDGMENTS

Eternal gratitude from A.C. goes to Jo Chamblin (Piglit) for loving support and help with the preparation of this paper. A.S. acknowledges financial support from the S.E.R.C., and A.C. is supported by N.S.F. Graduate Fellowship No. RCD-9255644.

- [1] D. Deutsch, Phys. Rev. **D44**, 3197 (1991).
- [2] H.D. Politzer, Phys. Rev. **D46**, 4470 (1992).
- [3] G.W. Gibbons and S.W. Hawking, Phys. Rev. Lett. **69**, 1719 (1992).
- [4] A. Chamblin and R. Penrose, Twistor Newsletter **34**, 13 (1992).
- [5] A. Chamblin, J. Geom. Phys. (to appear) (1993).
- [6] G.W. Gibbons and S.W. Hawking, Comm. Math. Phys. **148**, 345 (1992).